

A comparison of FQHE quasi-electron trial wave functions on the disk

M. Kasner and W. Apel
Physikalisch-Technische Bundesanstalt
Bundesallee 100
D-38116 Braunschweig, Germany

Abstract

The various proposals for FQHE quasi-electron trial wave functions are reconsidered. In a short-range model for the electronic interaction, the energy expectation values of four different trial wave functions are calculated at filling factor $\frac{1}{3}$ for up to ten electrons in the disk geometry. Jain's trial wave function displays the lowest energy expectation value.

PACS-Numbers: 73.20Dx, 73.50Jt

It is by now generally accepted that the fractional quantum Hall effect (FQHE) arises due to peculiar features of the multiparticle energy spectrum of interacting, spin-polarized electrons moving in two dimensions under the influence of a strong, perpendicular magnetic field [1]. These peculiarities appear at definite filling factors ν (electronic densities). Best understood are the ground states at filling factors $\nu = \frac{1}{q}$ (q - odd). Here, the Laughlin wave function [2] is not only a good trial wave function, but also an exact solution of a special short-range two-particle interaction [3]. Quasi-particles were introduced in order to describe the low-lying energy levels in systems with a few flux quanta off from $\nu = \frac{1}{q}$. They were also constructed in order to explain the occurrence of the other filling factors $\nu = \frac{p}{q}$ by the concept of a condensation of these quasi-particles, quite analogous to the condensation of the interacting electrons into the $\frac{1}{q}$ -state [4]. However, this picture is based on some prepositions. These are the dominance of the short-range contribution in the pair-interaction of the quasi-particles and the smallness of the interaction between these quasi-particles in comparison with the energy gap of the collective excitations of the parent state [5]. Also a microscopic derivation of the quantum mechanics of these quasi-particles from first principles is still missing.

The theory received renewed interest when a different approach towards an explanation of filling factors unequal $\frac{1}{q}$ was proposed by Jain [6], who gave an explicit prescription not only for constructing trial wave functions for any number of quasi-particles at $\nu = \frac{1}{q}$ but also for ground states and even the lowest excitations at all the other filling factors. The idea was to exploit the incompressibility of non-interacting electrons at integer filling factors. A multiplication with a symmetric Jastrow-factor then leads to a correlated wave function which has the desired filling factor. Finally, the wave function is projected onto the lowest Landau level $n = 0$, an almost unnecessary procedure, since after the multiplication with the Jastrow-factor, only a little weight remains in higher Landau levels [7]. This formulation was successfully used to explain multiparticle spectra on the sphere [8]. On the other hand, there are many quantitative results supporting the previous hierarchy scheme [9]. However, the relation between both approaches is still unclear.

In order to study the relative merits of these two approaches, we choose the simplest case where differences can appear. While the construction of the $\nu = \frac{1}{3}$ -quasi-hole leads in either case to the same trial wave function, the quasi-electron of the $\frac{1}{3}$ -state given by Jain differs from the quasi-electron proposed originally by Laughlin. It is not surprising that, in contrast to the case of the quasi-holes, there are many proposals for different quasi-electron trial wave functions [10, 11, 12].

The Laughlin wave function is quite a good trial wave function for various two-particle interactions, and this expresses the universality of the FQHE. Nevertheless, the characteristic features of the ground state can be destroyed, if the short-range contribution no longer dominates the interaction [5]. Therefore, and because the Laughlin wave function is an exact solution to this, we use the short-range model as a canonical model.

In a first attempt, we aim at the properties of the $\nu = \frac{1}{3}$ -quasi-electron. Here, we work in the disk geometry [13] which was originally used to formulate Laughlin's idea, although we are aware of some drawbacks in comparison with the spherical geometry.

Crucial for the quasi-electron construction in the hierarchical scheme is that all angular momentum components of the quasi-electron are degenerated in energy. On the sphere, this is a priori fulfilled because of the symmetry. For a system in the disk geometry with a finite particle number N , this is not satisfied, but in the thermodynamic limit the energies for all the components of a trial quasi-electron wave function should tend to the same limiting value. Deviations from the limit can be assigned to i) the quality of the proposed trial wave function and, of course, ii) to the influence of the finite size of the system. The finite size corrections of the ground state energy, e. g. are proportional to $\frac{1}{\sqrt{N}}$ on a disk and proportional to $\frac{1}{N}$ on a sphere. It is our aim to study the full dependence of the quasi-electron energies on the angular momentum for finite N . In previous studies of the disk geometry, the quasi-electron at the origin was considered corresponding to just a single angular momentum component.

The Hamiltonian to be investigated describes N spin-polarized electrons moving on a two-dimensional disk. The Hilbert space is restricted to the lowest Landau level and thus contains just an interaction term because we do not include a background in our model. It reads:

$$H = \frac{1}{2} \sum_{m_1, m_2, m_3, m_4=0}^{m_{max}} W_{m_1 m_2 m_3 m_4} c_{m_1}^+ c_{m_2}^+ c_{m_3} c_{m_4}, \quad (1)$$

where c_m^+ creates an electron state with angular momentum m in the lowest Landau level ($m = 0, 1, \dots$). The matrix elements $W_{m_1 m_2 m_3 m_4}$ are specified such that only the pseudo-potential coefficient $V_1 = \frac{\sqrt{\pi}}{4}$ is nonzero [5], i. e. only electrons with relative angular momentum one repel each other.

The calculations are performed for the quasi-electrons of the $\frac{1}{3}$ -state. Because the filling factor is defined by the relation $\nu = \frac{N-1}{m_{max}} = \frac{N-1}{N_\Phi-1}$, where m_{max} is the maximum single particle angular momentum and limits the area of the disk and where N_Φ is the number of flux-quanta through this disk, the relation $m_{max} = 3(N-1)$ holds for the stable state at $\nu = \frac{1}{3}$. If we increase the density of the system by keeping N fixed while decreasing the area by the area of just one flux quantum, we create a quasi-electron. Thus, for a one-quasi-electron trial wave function at $\nu = \frac{1}{3}$ we must have

$$m_{max} = 3(N-1) - 1. \quad (2)$$

In this work we report calculations of the energy expectation values of the angular momentum components for four different quasi-electron wave functions proposed in the literature. This is done for up to $N = 10$ particles.

The quasi-electron wave functions have total angular momenta M reaching

from $M^* - N$ to $M^* - 1$, where $M^* = \frac{q}{2}N(N-1)$ is the total angular momentum of the $\frac{1}{q}$ -Laughlin state [13]. The states with higher angular momenta are attributed to the edge of the system [14].

The first proposal for a quasi-electron wave functions is due to Laughlin [2]:

$$\Psi_{\frac{1}{q}}^{(+,z_0)} = \prod_{i=1}^N e^{-\frac{1}{4}|z_i|^2} (2\partial_{z_i} - z_0^*) \prod_{k<l}^N (z_k - z_l)^q. \quad (3)$$

This trial wave function can either be viewed as a quasi-electron located at a position z_0 or it can be expanded with respect to z_0 in order to get components, which are eigenfunctions of the total angular momentum varying from $M^* - N$ to M^* (the state with $M^* - 1$ is missing). All components of the Laughlin quasi-electron wave function, except the one at angular momentum $M^* - N$, contain Slater determinants with single particle angular momentum $3(N-1)$ and thus violate the constraint (2). Only in the component at $M^* - N$ all derivatives act on the polynomial part in (3) and, thus, here the constraint (2) is met. Therefore, we have also studied a modified Laughlin quasi-electron trial wave function $\tilde{\Psi}_{\frac{1}{q}}^{(+,z_0)}$, in which all Slater determinants containing single particle angular momenta $m > m_{max}$ are excluded. This wave function has the correct filling factor, but one gets higher energy.

The next quasi-electron trial wave function was proposed by MacDonald and Girvin [10]. It is written in occupation number representation:

$$|\phi_{\frac{1}{q}}^{(+)}\rangle = \hat{d}_{\infty} \dots \hat{d}_1 (1 - \hat{n}_0) |\Psi_{\frac{1}{q}}\rangle. \quad (4)$$

The operator defined as $\hat{d}_m = c_{m-1}^+ c_m + 1 - n_m$ decreases all one-particle angular momenta by one. All states containing zero one-particle angular momentum are projected out by $(1 - \hat{n}_0)$. Unfortunately, one obtains only a single component (with $M = M^* - N$).

As already mentioned, the third trial function is due to Jain [6]. Its component with angular momentum $M^* - N + m + 1$ ($m = -1, 0, \dots, N-2$) is given by

$$\chi_{\frac{1}{3}}^{(+,M^*-N+m+1)} = \mathcal{P}_0 \prod_{i<j}^N (z_i - z_j)^2 \begin{vmatrix} z_1^m |z_1|^2 & \dots & z_N^m |z_N|^2 \\ 1 & \dots & 1 \\ z_1 & \dots & z_N \\ \vdots & \ddots & \vdots \\ z_1^{N-2} & \dots & z_N^{N-2} \end{vmatrix} \prod_{i=1}^N e^{-\frac{1}{4}|z_i|^2}. \quad (5)$$

$\mathcal{P}_0$ is the above mentioned projector onto the lowest Landau level.

A fourth proposal based on the projection of the trial wave function from the sphere onto the plane was made by us [12]. This method generates $N + 1$ angular momentum components resulting from the expansion of

$$\Phi_{\frac{1}{q}}^{(+,z_0)} = \prod_{i=1}^N ((N-1)q - z_i \partial_{z_i} + z_0 \partial_{z_i}) \Psi_{\frac{1}{q}} \quad (6)$$

with respect to z_0 . In (6), the derivatives act as usual only on the polynomial part of $\Psi_{\frac{1}{q}}$. All the components of (6) have the correct filling factor (condition (2) is fulfilled).

Expanding the components of all trial wave functions (3)-(6) in Slater determinants, we calculated numerically exact the coefficients for particle numbers up to $N = 10$. From these coefficients, then the energy expectation values were determined for $\Psi_{\frac{1}{3}}^{(+)}$, $\tilde{\Psi}_{\frac{1}{3}}^{(+)}$, $\phi_{\frac{1}{3}}^{(+)}$, $\chi_{\frac{1}{3}}^{(+)}$ and $\Phi_{\frac{1}{3}}^{(+)}$. The results are shown for $N = 8, 9, 10$ in Table 1a, b and c, respectively. The data are pictured in Fig. 1 for $N=8$.

All data for $N = 8, 9, 10$ show a very similar behaviour with respect to their absolute values as well as to their dependence on the angular momentum M . In the case of 8 particles we added, for the sake of comparison, the lowest energy for each M resulting from the exact numerical diagonalization of H , where the maximum single particle angular momentum is taken to be $m_{max} = 3(N-1) - 1$. It is quite obvious that the exact energies do not depend strongly on M and this supports the picture of degeneracy for $M = M^* - N$ to $M^* - 1$. The absolute values of the data are near the quasi-electron energy at $\nu = \frac{1}{3}$: 0.1905, which was estimated by extrapolation of the finite-size data from calculations on the sphere, see Table 2 in [15]. The differences between the energies of all the trial wave functions and the exact energies increase with increasing M .

For all N and all M -components, Jain's quasi-electron wave function is most favorable. The M -dependences of our and Jain's proposal parallel each other, but our data are always larger. The difference between the energies of Laughlin's proposal (modified to meet (2), second column in Table 1) and the ones of Jain's proposal increases with increasing M for all N . We include in the Tables also the energy expectation values of Laughlin's original proposal (column 1). These energy expectation values differ only about 2% from the ones of the modified Laughlin trial wave function which meets the condition (2). Therefore, while the objection that Laughlin's original quasi-electron does not show the correct filling factor (m_{max}) is certainly true, its energies are not so much affected by the violation of the condition (2).

In the last row of the Tables, we include the energies for the wave functions with angular momentum M^* . These data are well separated from those of the other angular momenta, e. g. the expectation value for $N = 10$ and $M = 135$ in Table 1c is much lower than all the other ones. Thus, we identify only the N lowest M -components with the quasi-electron state, cf. [13].

For $N = 8$, we calculate the mean of the N quasi-electron components from Laughlin's ($N - 1$ values), Jain's and our trial wave functions and from the exact data. We find 0.3554, 0.3040, 0.3240 and 0.1942, respectively. Thus, the relative deviations of the data of the three trial wave functions from the exact ones are 83%, 57% and 66%. Jain's proposal is closest to the exact value, but it is not as good as the corresponding proposal in the spherical geometry. In general, all the energies of the trial wave functions are still rather high so that there is still room for improvement.

In conclusion, the quasi-electron on the disk can be identified in the energy spectrum. There are N angular momentum components. The various proposals for trial wave functions show rather different behaviour and are of quite different quality. Our approach of studying all the angular momentum components of the quasi-electron in the disk geometry leads to a very detailed picture of the behaviour of the various proposals for trial wave functions and allows for a clear distinction of the relative merits of these.

We are grateful for stimulating discussions with J. K. Jain, A. H. MacDonald and W. Weller. We acknowledge partial support by the Deutsche Forschungsgemeinschaft via grant Ap47/1-1 and by the European Community within the SCIENCE-project, grant SCC*-CT90-0020.

Table 1a

M	$\Psi_{\frac{1}{3}}^{(+)}$	$\tilde{\Psi}_{\frac{1}{3}}^{(+)}$	$\phi_{\frac{1}{3}}^{(+)}$	$\chi_{\frac{1}{3}}^{(+)}$	$\Phi_{\frac{1}{3}}^{(+)}$	num. diagon.
76	0.2413	0.2413	0.2362	0.2033	0.2413	0.1903
77	0.2191	0.2199	—	0.1932	0.2200	0.1828
78	0.2467	0.2491	—	0.2246	0.2474	0.1982
79	0.3141	0.3183	—	0.2799	0.2980	0.1964
80	0.3966	0.4022	—	0.3319	0.3445	0.1968
81	0.4879	0.4954	—	0.3737	0.3835	0.1955
82	0.5822	0.5920	—	0.4034	0.4159	0.1954
83	—	—	—	0.4216	0.4411	0.1983
84	0	0.0137	—	0.0169	0.0896	0.0132

Table 1b

M	$\Psi_{\frac{1}{3}}^{(+)}$	$\tilde{\Psi}_{\frac{1}{3}}^{(+)}$	$\phi_{\frac{1}{3}}^{(+)}$	$\chi_{\frac{1}{3}}^{(+)}$	$\Phi_{\frac{1}{3}}^{(+)}$
99	0.2467	0.2467	0.2438	0.2065	0.2467
100	0.2232	0.2239	—	0.1926	0.2240
101	0.2284	0.2302	—	0.2047	0.2314
102	0.2715	0.2747	—	0.2447	0.2688
103	0.3379	0.3423	—	0.2943	0.3141
104	0.4142	0.4198	—	0.3392	0.3546
105	0.5000	0.5072	—	0.3762	0.3897
106	0.5857	0.5947	—	0.4028	0.4192
107	—	—	—	0.4202	0.4437
108	0	0.0119	—	0.0148	0.0926

Table 1c

M	$\Psi_{\frac{1}{3}}^{(+)}$	$\tilde{\Psi}_{\frac{1}{3}}^{(+)}$	$\phi_{\frac{1}{3}}^{(+)}$	$\chi_{\frac{1}{3}}^{(+)}$	$\Phi_{\frac{1}{3}}^{(+)}$
125	0.2495	0.2495	0.2463	0.2082	0.2495
126	0.2309	0.2316	—	0.1960	0.2316
127	0.2221	0.2235	—	0.1954	0.2255
128	0.2440	0.2464	—	0.2191	0.2471
129	0.2931	0.2966	—	0.2603	0.2861
130	0.3566	0.3610	—	0.3047	0.3266
131	0.4290	0.4345	—	0.3449	0.3633
132	0.5098	0.5164	—	0.3780	0.3953
133	0.5890	0.5972	—	0.4024	0.4227
134	—	—	—	0.4189	0.4462
135	0	0.0105	—	0.0132	0.0956

Captions:

Table 1(a-c): The energy expectation values for the trial wave functions (3)-(6) for $N = 8$ (1a), $N = 9$ (1b) and $N = 10$ (1c) particles in dependence on angular momentum M . For $N = 8$, also the results of the exact diagonalization are given.

Fig. 1: The energy expectation values of Laughlin's (dashed line), Jain's (solid line) and our (broken line) quasi-electron trial wave function plotted against angular momentum M for $N = 8$ particles. The dotted lines correspond to the exact values. The data are given in Table 1a.

References

- [1] A. H. MacDonald, *Helv. Phys. Acta* **65**, 133 (1992).
- [2] R. B. Laughlin, *Phys. Rev. Lett.* **50**, 1395 (1983).
- [3] F. D. M. Haldane, *Phys. Rev. Lett.* **51**, 605 (1983).
- [4] B. I. Halperin, *Phys. Rev. Lett.* **52**, 1583, 2390 (E) (1984).
- [5] F. D. M. Haldane, in *The Fractional Quantum Hall Effect*, edited by R. E. Prange and S. M. Girvin (Springer-Verlag, New York, 1990).
- [6] J. K. Jain, *Phys. Rev. B* **40**, 8079 (1989).
- [7] N. Trivedi and J. K. Jain, *Mod. Phys. Lett. B* **5**, 503 (1991).
- [8] G. Dev and J. K. Jain, *Phys. Rev. Lett.* **69**, 2843 (1992).
- [9] P. Béran and R. Morf, *Phys. Rev. B* **43**, 12654 (1991).
- [10] A. H. MacDonald and S. M. Girvin, *Phys. Rev. B* **33**, 4414 (1986).
- [11] R. Morf and B. I. Halperin, *Phys. Rev. B* **33**, 2221 (1986).
- [12] M. Kasner and W. Apel, *Europhys. Lett.* **20**, 715 (1992).
- [13] M. Kasner and W. Apel, to be published.
- [14] A. H. MacDonald, *Phys. Rev. Lett.* **64**, 220 (1990).
- [15] C. Gros and A. H. MacDonald, *Phys. Rev. B* **42**, 9514 (1990).